


Composition and size of dust grain.

Radiation field intensity and spectrum


Composition and size of dust grain.



Optical properties of grain


Radiation field intensity and spectrum

Composition and size of dust grain.




Optical properties of grain


Radiation field intensity and spectrum



Grain temperature pdf




For a given radiation field intensity, larger grains can be in equilibrium, but small grains may not be.




U = radiation field intensity scaled to MMP Solar neighborhood field

Composition and size of dust grain.





Optical properties of grain

Radiation field intensity and spectrum




Grain temperature pdf






Emission spectrum





grains in equilibrium with the radiation field, T~U^{1/6}

stochastically heated grains

Draine & Li 2007










What happens if you change the radiation field spectrum?

Matters mostly for small stochastically heated grains, where the average photon energy determines peak T.






What happens with a distribution of radiation field intensities?



Distribution of U (equivalent to distribution of T):

- broadens far-IR peak
- changes long wavelength slope
- increases mid-IR for similar MBB peak

What happens with a distribution of radiation field intensities?





Distribution of U (equivalent to distribution of T):

- broadens far-IR peak - changes long wavelength slope - increases mid-IR for similar MBB peak

Common practice: power-law distribution for dM/dU with delta function at Umin


> (Dale+2001, Draine & Li 2007, Aniano+2020)











IRAS - 0.57 m































Spitzer - 0.85 m







Herschel - 3.5m











Credit: NASA, ESA, CSA, STScl, Janice Lee (STScl), Thomas Williams (Oxford), PHANGS Team




Credit: NASA, ESA, CSA, STScl, Janice Lee (STScl), Thomas Williams (Oxford), PHANGS Team




Things we would like to learn from dust emission:

- How does dust mass vary within and between galaxies?
- What controls the small carbonaceous grain (PAH) fraction?
- Do the properties of PAHs vary throughout the ISM?
- Can we use dust emission as a tracer of star formation and/or gas?



Dust Mass Scaling Relations



star formation "main sequence"

M_{star} - Z

mass-metallicity relation

M_{star} - M_{gas}
gas fraction

Mgas - SFR

Kennicutt-Schmidt relation



Dust-to-metals is the link that connects dust mass scaling laws to galaxy properties.

FR Scaling Laws

M_{dust} - M_{star}

M_{dust} - SFR

M_{dust} - M_{gas}

Just Sca

Dust Mass Scaling Relations



star formation "main sequence"

M_{star} - Z

mass-metallicity relation

M_{star} - M_{gas}
gas fraction

Mgas - SFR

Kennicutt-Schmidt relation



Dust-to-metals is the link that connects dust mass scaling laws to galaxy properties.

FR Scaling Laws

8

Mstar (

~all nearby galaxies

~1000 nearby galaxies


Dust Sca

<100 nearby galaxies

M_{dust} - M_{star}


M_{dust} - SFR

M_{dust} - M_{gas}




 M_{dust}/M_{star} trend with sSFR reflects combination of gas fraction, Z, and D/M.





- Difference between absorption (depletion) and emission based D/M behavior.
 - Big scatter in galaxy-integrated emission values at fixed Z.
 - Sharp drop in emission values at low Z.
- Is this assumed far-IR emissivity issues?
- Is this because of smearing out metallicity & DGR gradients in galaxies vs lines-of-sight?

Resolved Dust-to-Metals Ratio





Emission-based D/M highly tied to assumed dust opacity, X_{CO}.

We have all of the far-IR we're going to get for a while.

There are ~1000 resolved galaxies with far-IR observations from Herschel.

https://irsa.ipac.caltech.edu/data/Herschel/z0MGS_Dust/overview.html






All Herschel maps and dust SED fit results available at IRSA (Chastenet+2025)

See also DustPedia database (Davies+2017,2019; Clark+2018,2019)


What controls the PAH fraction?

PAH Fraction = fraction of dust mass in the form of carbonaceous grains with <10³ C atoms.



What controls the PAH fraction?

PAH Fraction = fraction of dust mass in the form of carbonaceous grains with <10³ C atoms.




From peak of modified blackbody, get U (or T) and M_{dust}.

With known U and M_{dust}, scaling of mid-IR gives PAH mass fraction.

What controls the PAH fraction?

Mid-IR JWST-based tracer RPAH tracks qPAH at far higher angular resolution.



Based on mid-IR tracers of PAH fraction with Spitzer: Engelbracht+2005, 2008, Marble+2010

What controls the PAH fraction?

Two main behaviors:

1) Destruction of PAHs in ionized gas.


2) Deficit of PAHs at low metallicity.

The lack of PAHs in regions of ionized gas has been clear for many years with observations from IRAS, ISO, and Spitzer in Milky Way HII regions.


Cesarsky+1996, Kassis+2006, Povich+2007

With sufficient angular resolution, can see this in other galaxies.

 $3-500 \, \mu m \, dust \, fitting -> 10 \, pc \, map \, of \, q_{PAH}$ in Magellanic Clouds.





~5-10 pc resolution now possible in galaxies out to ~10 Mpc with JWST MIRI




Chastenet et al. 2023a, Sutter et al. 2024


~5-10 pc resolution now possible in galaxies out to ~10 Mpc with JWST MIRI









PAH destruction in HII regions matters on the scale of entire galaxies!







Observations with ISO and Spitzer have shown a deficit of PAH emission from lowmetallicity galaxies.

Lots of evidence for PAH-Z deficiency:

Madden et al. 2000

Galliano et al. 2003, 2005

Engelbracht et al. 2005


Madden et al. 2006

Wu et al. 2006

Draine et al. 2007

Remy-Ruyer et al. 2015

etc., etc.



Observations with ISO and Spitzer have shown a deficit of PAH emission from lowmetallicity galaxies.

Lots of evidence for PAH-Z deficiency:

Madden et al. 2000

Galliano et al. 2003, 2005

Engelbracht et al. 2005


Madden et al. 2006


Wu et al. 2006


Draine et al. 2007

Remy-Ruyer et al. 2015

etc., etc.








Somewhere between the SMC & LMC's metallicity there is a dramatic change in PAH abundance.

WHY?


Aniano et al. 2020 (also Draine et al. 2007)

Chastenet et al. 2019


Part of the story may be a low filling factor of regions with high qPAH.




Part of the story may be a low filling factor of regions with high qPAH.



Part of the story may be a low filling factor of regions with high qPAH.






Tarantino+2026 (in prep)

Sandstrom+2010


Another (related) part may be due to inhibited growth of PAHs at low Z.



Short λ features increase in importance at low Z, suggests smaller PAHs. Destruction generally leads to larger PAHs -> "inhibited growth" explanation.


Sandstrom+2012, Whitcomb+2024, 2025, Tarantino+2026 (in prep)

Another (related) part may be due to inhibited growth of PAHs at low Z.



Short λ features increase in importance at low Z, suggests smaller PAHs. Destruction generally leads to larger PAHs -> "inhibited growth" explanation.


Sandstrom+2012, Whitcomb+2024, 2025, Tarantino+2026 (in prep)



particularly dramatic - Lai+2025 II Zw 40

Do properties of PAHs vary throughout the ISM?


Ratios of major mid-IR features trace size, charge, and radiation field hardness.



Allamandola et al. 1999 (reproduced in Tielens 2008)

Do properties of PAHs vary throughout the ISM?


Ratios of major mid-IR features trace size, charge, and radiation field hardness.



Allamandola et al. 1999 (reproduced in Tielens 2008)


Draine & Li 2007, Draine et al. 2021

How important is the radiation field spectrum?




One of the strongest trends is between optical line ratios and PAH band ratios.

How important is the radiation field spectrum?



Ionized gas & PAHs see different parts of the radiation field.


How important is the radiation field spectrum?



Observed trends are consistent with ~constant size & charge and varying radiation field spectrum!


See also poster by Deb Pathak!

Yes, but it is always a bit of both!



1) Dust is well-mixed with gas to first order.


Yes, but it is always a bit of both!



- 1) Dust is well-mixed with gas to first order.
- 2) Depending on λ , emission depends either linearly (e.g. 7.7) or non-linearly (e.g. 70) with U.

Draine & Li 2007


Yes, but it is always a bit of both!



- 1) Dust is well-mixed with gas to first order.
- 2) Depending on λ , emission depends either linearly (e.g. 7.7) or non-linearly (e.g. 70) with U.
 - 3) U increases rapidly near massive star forming regions.


Draine & Li 2007

Yes, but it is always a bit of both!




- 1) Dust is well-mixed with gas to first order.
- 2) Depending on λ , emission depends either linearly (e.g. 7.7) or non-linearly (e.g. 70) with U.
 - 3) U increases rapidly near massive star forming regions.

Yes, but it is always a bit of both!



- 1) Dust is well-mixed with gas to first order.
- 2) Depending on λ , emission depends either linearly (e.g. 7.7) or non-linearly (e.g. 70) with U.
 - 3) U increases rapidly near massive star forming regions.
 - 4) U includes contributions from older stellar pops.







Correlation of ~kpc SFR and CO with mid- to far-IR dust emission in SINGS.


Contribution from hot dust in equilibrium with high U & non-linear U dependence leads to better correlation with SFR.

Stochastic emission and cold equilibrium dust emission better correlate wth gas.




In H_2 dominated regions of galaxies $(\Sigma_{gas} \sim \Sigma_{H2})$, strong, ~linear correlation of PAH emission with CO.







PAH emission with JWST is the highest available resolution & sensitivity ISM tracer. May be very useful as a ~total gas surface density tracer for comparison to simulations. image courtesy of Matt Orr - FIRE + PHANGS



Many exciting possibilities for using PAH emission as a gas tracer.



Thilker+2023, Meidt+2023




What comes next?



Critical to have access to far-IR.

SPHEREX



6" resolution near-IR spectral map of every nearby galaxy!




Conclusions


- Dust-to-metals links galaxy scaling relationships to dust. Need to understand the discrepancies between emission and absorption based measurements.
- PAHs are destroyed in ionized gas, and this can impact global PAH fractions in high SFR conditions.
- PAH fraction drops with metallicity perhaps due to a combination of filling factor and more fragile, easily destroyed PAHs.
- The degree to which PAH band ratios reflect size/charge or radiation field variations is still tbd.
- Dust emission traces both gas and star formation depending on the circumstances.

Spitzer IRS Mapping Legacy Archive

Donnelly et al. 2025 (in prep) - delivery to IRSA later this year




Delivering of all IRS low resolution spectral mapping cubes.





Spitzer IRS Mapping Legacy Archive

Donnelly et al. 2025 (in prep) - delivery to IRSA later this year



Delivering of all IRS low resolution spectral mapping cubes.





https://www.aromaticuniverse.space/